

SOCIAL CONTEXT MATTERS

Facilitated Differential Pain-related Fear Learning in a Threatening Social Context

Kai Karos¹, Ann Meulders^{1,2} and Johan W.S. Vlaeyen^{1,3}

¹ Department of Psychology, Research Group on Health Psychology, University of Leuven, Belgium

² Center for Excellence on Generalization Research in Health and Psychopathology, University of Leuven, Belgium

³ Faculty of Psychology, Department of Clinical Psychological Science, Maastricht University, The Netherlands

aps

ASSOCIATION FOR PSYCHOLOGICAL SCIENCE

KU LEUVEN OGP

Introduction

- Pain-related fear** plays an important role in the development and maintenance of **chronic pain**, and is a major focus in current treatment for chronic pain¹
- Factors facilitating the development of perpetual maladaptive fear are unknown**, making the prevention of chronic pain challenging.
- A novel approach: The study of pain in relation to social contextual factors. **Social context** modulates the appraisal, interpretation, and experience of pain², which might make it a feasible target for prevention.

Aim: The current study investigated the effects of a threatening and a safe social context on pain-related fear using an established differential voluntary fear conditioning paradigm (i.e., the "Voluntary Joystick Movement (VJM) Paradigm")³.

Hypotheses: A threatening social context compared to a safe social context leads to:

- Enhanced (i.e., faster / stronger) acquisition of pain-related fear
- Slower extinction of pain-related fear
- Increased pain intensity ratings
- Increased contextual fear
 - Increased baseline startle responses
 - Increased startle responses during the inter trial intervals (ITI)
 - Slower reaction times

Methods

Stimuli

Unconditioned Stimulus (US)



Pain-US
Electrocutaneous stimulation
(M = 33.79 mA;
SD = 18.35)

Conditioned Stimulus (CS)



Proprioceptive Stimuli
left/right movements


Social Context Manipulation



Facial stimuli
Angry (threat group) or happy (safe group) facial stimuli⁴

Participants & Design

N = 42 healthy participants (12 ♂)
Age: 17-29 years old
2 groups: Threat (n=21) vs. Safe (n=21)



During acquisition a shock followed the left movement (CS+), but not the right movement (CS-).

Measures

Self-report measures (0-10):

- Pain expectancy
- Pain-related fear
- Pain intensity
- Threat of pain


Behavioral measures:

- Movement-onset latency
- Response duration




Psychophysiological measures:


- Eyeblink startle modulation elicited by startle probes (white noise, 100 dB)



Design



Results



- CS+ vs. CS- (ACQ3) Threat > Safe (Pain Expectancy, Pain-related fear, Startle Amplitude) ✓
- Threat = Safe; Extinction not complete at EXT3 ✗
- Threat = Safe; no differences in pain intensity ratings ✗
- Increased contextual fear ✓
 - Threat > Safe (Baseline Startle) ✓
 - Threat > Safe (EXT3, ITI) p=.07 ✗
 - Threat < Safe (EXT) ✓

Note: * = p<.05 | ** = p<.01

Conclusion

A threatening social context

- leads to contextual fear** - Behavioral "freezing" and increased physical arousal⁵
- facilitates differential fear learning** - Evolutionary "preparedness" to distinguish between threat and safety⁶ / threatening context motivates efficient differential learning
- Does not affect pain intensity ratings**
- Does not modulate extinction of pain-related fear**

Summary: Exposure to a threatening social context affects the perception of pain by modulating pain-related fear and contextual fear. Increased clinical and scientific attention to the effects of social context in acute and chronic pain conditions is warranted.

Correspondence to:

Kai Karos

Kai.karos@ppw.kuleuven.be

<http://ppw.kuleuven.be/home/english/research/ogp>

References:

- Vlaeyen, J. W. S., & Linton, S. J. (2012). Fear-avoidance model of chronic musculoskeletal pain: 12 years on. *Pain*, 153(1), 11-19. doi:10.1016/j.pain.2011.12.009
- Hadjistavropoulos, T., Craig, K. D., Duck, S., Cano, A., Goubert, L., Jackson, P. L., ... Fitzgerald, T. D. (2011). A biopsychosocial formulation of pain communication. *Psychological Bulletin*, 137(6), 910-939. doi:10.1037/a0023876
- Meulders, A., Vansteenwegen, D., & Vlaeyen, J. W. S. (2011). The acquisition of fear of movement-related pain and associative learning: a novel pain-relevant human fear conditioning paradigm. *Pain*, 152(11), 2440-9. doi:10.1016/j.pain.2011.05.015
- Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., ... Nelson, C. (2009). The NimStim set of facial expressions: judgments from untrained research participants. *Psychiatry Research*, 168(3), 242-9. doi:10.1016/j.psychres.2008.05.006
- Roelofs, K., Hagenaars, M. A., & Sijns, J. (2010). Facing freeze: social threat induces bodily freeze in humans. *Psychological Science*, 21(11), 1575-81. doi:10.1177/0956797610384746
- Östman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: Toward an evolved module of fear and fear learning. *Psychological Review*, 108(3), 483-502. doi:10.1037/0033-295X.108.3.483

